

INDEX TO VOLUME 113

New scientific names are in bold face

- Abundance 187–200
Acadia National Park 260–279
Acrosiphonia arcta (Codiolales, Chlorophyta) from environmentally different habitats in Nova Scotia, Canada. Morphological variation in, 87–105
Acrosiphonia arcta, *A. sonderi*, *A. spinescens*, 87–105
Aedo, Carlos and Patrick J. Alexander. *Geranium dodecatheoides*, a new species from New Mexico, U.S.A. 252–259
Aegagropilous *Gracilaria* 300–350
Alexander, Patrick J. 252–259
Alix, Mitchell S. 32–46
Allegheny High Plateau 47–63
Angelo, Ray and David E. Boufford. Atlas of the flora of New England: Caryophyllidae. 419–513; maps 420, 447–512
ANNOUNCEMENTS:
A New Feature in *Rhodora* 123, 234, 364
Merritt Lyndon Fernald Award 363
NEBC Graduate Student Research Award 362
Aster 64–86
Asteraceae 64–86
Atlantic Coastal Plain 32–46
Atlas 419–513
Atlas of the flora of New England: Caryophyllidae. 419–513; maps 420, 447–512
Baltic Sea (northern Europe) 300–350
Bart, David, Matt Simon, Quentin Carpenter, and Stephanie Graham. Historical land use and plant-community variability in a Wisconsin calcareous fen. 160–186; species list 183–186
Benthic “marine” algae in Bras d’Or Lake, Nova Scotia with five other Northwest Atlantic embayments and the Baltic Sea in northern Europe. A floristic comparison of, 300–350; species list for Bras d’Or Lake 334–350
Biogeography 260–279
Biomonitoring 1–31
Bladderwort 32–46
BOOK REVIEWS:
Botanic Gardens: Modern-Day Arks 225–226
Plants of Acadia National Park, The 353–354
Plants of St. Lawrence County, NY: An Annotated Checklist of Vascular Flora 106–108
Serpentine: The Evolution and Ecology of a Model System 523–526
William Bartram, the Search for Nature’s Design: Selected Art, Letters, and Unpublished Writings 109–112
Borkowsky, Christie L. 201–219
Boufford, David E. 419–513
Brackish water 300–350
Bras d’Or Lake, Nova Scotia 300–350; species list 334–350
Brooksville, Maine, U.S.A. 1–31
Budnick, Karen E. 201–219
Buhrstone Hills, Mississippi 365–418
Calcareous fens 160–186
Cape Breton Island 300–350
Carnivorous plants 235–251
Carpenter, Quentin 160–186
Caryophyllidae 419–513
Chlorophyta 87–105
Ciccotelli, Brett, Tanner B. Harris, Bruce Connery, and Nishanta Rajakaruna. A preliminary study of the vegetation of vernal pools of Acadia National Park, Maine, U.S.A. 260–279; species list 267–273
Circumboreal 187–200

- Clayden, Stephen R. 1-31
Composition and structure of riparian forest dominated by *Tsuga canadensis*: Contrasts between old-growth and second-growth stands in northwestern Pennsylvania. 47-63
Conioselinum chinense over-reported in eastern Massachusetts due to confusion with *Peucedanum palustre*. 516-518 (The Botanists' Corner)
Connery, Bruce 260-279
Connolly, Bryan A. *Conioselinum chinense* over-reported in eastern Massachusetts due to confusion with *Peucedanum palustre*. 516-518 (The Botanists' Corner)
Conservation 1-31
Cook Forest State Park 47-63
Cook, Rachael E. and Charles E. Williams. Composition and structure of riparian forest dominated by *Tsuga canadensis*: Contrasts between old-growth and second-growth stands in northwestern Pennsylvania. 47-63
Copper- and zinc-enriched superfund site in Brooksville, Maine, U.S.A. A, 1-31
Cyperaceae 351-352
Dawes, Clinton J. 300-350
Dibble, Alison C. 1-31
Distribution 419-513
Disturbance 187-200
Dodecatheon 280-299
Doffitt, Chris 365-418
Doughty, E. D. 235-251
Dwarf scouring-rush 187-200
Eastern hemlock 47-63
Edaphic ecology 1-31
Edaphic islands 260-279
Ellison, A. M. 235-251
Endangered species 32-46; 516-518
Endemism 1-31
Environmental factors influencing the local abundance of *Equisetum scirpoideum* near the southern extent of its range. 187-200
Environmental pollution 1-31
Ephemeral wetlands 260-279
Equisetum scirpoideum 187-200
Equisetum scirpoideum near the southern extent of its range. Environmental factors influencing the local abundance of. 187-200
Esselman, Elizabeth J. 280-299
Estuarine 87-105; 300-350
Extremophiles 1-31
Flora (New England) 419-513
Floral visitor, insect 64-86
Floristic comparison of benthic "marine" algae in Bras D'Or Lake, Nova Scotia with five other Northwest Atlantic embayments and the Baltic Sea in northern Europe. A, 300-350; species list for Bras D'Or Lake 334-350
Floristic comparisons 300-350
Fruit and seed characters help distinguish southern Illinois *Dodecatheon* (Primulaceae) species and highlight unusual intergrading populations. 280-299
Gentianaceae 514-515
Geraniaceae 252-259
Geranium 252-259
Geranium dodecatheoides, a new species from New Mexico, U.S.A. 252-259
Graham, Stephanie 160-186
Habitat 187-200
Habitat conservation 260-279
Harris, Tanner B. 1-31; 260-279
Hemiparasitic plants 125-159
Hill, Jovonn 365-418
Historical land use and plant-community variability in a Wisconsin calcareous fen. 160-186; species list 183-186
Historical notes and new records for the rare Atlantic Coastal Plain species *Utricularia resupinata* (Lentibulariaceae) in Indiana. 32-46

- Horsburgh, M., J. C. Semple, and P. G. Kevan. Relative pollinator effectiveness of insect floral visitors to two sympatric species of wild aster: *Symphyotrichum lanceolatum* (Willd.) Nesom and *S. lateriflorum* (L.) Löve & Löve (Asteraceae: Astereae). 64–86
- Host plants 125–159
- Hoy, Joann. A new native plant for New Hampshire, *Trichophorum planifolium* (Cyperaceae). 351–352 (The Botanists' Corner)
- Hybridization 280–299
- Illinois, southern 280–299
- IN MEMORIAM:
- Leslie J. Mehrhoff 1950–2010 519–522
- Indiana flora 32–46
- Insular ecology 260–279
- Intergrading populations, *Dodecatheon* 280–299
- Invasive species 160–186
- Isolated wetlands 260–279
- Jones, Michael T. and William A. Patterson III. Environmental factors influencing the local abundance of *Equisetum scirpoides* near the southern extent of its range. 187–200
- Kevan, P. G. 64–86
- Lamiaceae 220–224
- Land-use history 160–186
- Lauderdale County, Mississippi 365–418
- Lentibulariaceae 32–46
- Lichen-metal relations 1–31
- Lichens of the Callahan Mine, a copper- and zinc-enriched superfund site in Brooksville, Maine, U.S.A. 1–31; species list 17–31
- Maine 1–31; 260–279
- Majure, Lucas C., Jovonn Hill, Chris Doffitt, and Terrence C. Majure. The vascular flora of Lauderdale County, Mississippi, U.S.A. 365–418; species list 388–418
- Majure, Terrence C. 365–418
- Manitoba 201–219
- Massachusetts 187–200; 516–518
- Mathieson, Arthur C. and Clinton J. Dawes. A floristic comparison of benthic "marine" algae in Bras D'Or Lake, Nova Scotia with five other Northwest Atlantic embayments and the Baltic Sea in northern Europe. 300–350; species list for Bras d'Or Lake 334–350
- Mississippi 365–418
- Morphological variability 87–105
- Morphological variation in *Acrosiphonia arcta* (Codiolales, Chlorophyta) from environmentally different habitats in Nova Scotia, Canada. 87–105
- Morphology 235–251
- Morphometrics 280–299
- Namestnik, Scott A. 32–46
- Naturalized species, *Salvia glutinosa* in southeastern New York 220–224
- Ne'man, G. 235–251
- Ne'man, R. 235–251
- NEBC Meeting News 114–122, 228–233, 356–361, 528–531
- Nectar 201–219
- Nectar sugar concentration and nectar quantity in the Western Prairie Fringed Orchid, *Platanthera prae-clara* (Orchidaceae). 201–219
- New Books 113, 227, 355, 527
- New England 187–200; 419–513
- New Hampshire 351–352
- New Mexico 252–259
- New native plant for New Hampshire, *Trichophorum planifolium* (Cyperaceae). A. 351–352 (The Botanists' Corner)
- New records 32–46; 351–352
- New species from New Mexico, U.S.A. *Geranium dodecatheoides* 252–259
- New York, *Salvia glutinosa* 220–224
- Non-native plants 125–159; 220–224

- Northwest Atlantic** 87–105
Northwest Atlantic embayments 300–350
Nova Scotia 87–105; 300–350
- Oberle, Brad and Elizabeth J. Esselman.** Fruit and seed characters help distinguish southern Illinois *Dodecatheon* (Primulaceae) species and highlight unusual intergrading populations. 280–299
- Olday, Fred C.** 1–31
- Old-growth and second-growth stands in northwestern Pennsylvania** 47–63
- Old-growth forest** 47–63
- Ontario, *Sympotrichum* floral visitors** 64–86
- Open coastal** 87–105
- Orchidaceae** 201–219
- Orobanchaceae** 125–159
- Oswald, W. W., E. D. Doughty, G. Ne'eman, R. Ne'eman, and A. M. Ellison.** Pollen morphology and its relationship to taxonomy of the genus *Sarracenia* (Sarraceniaceae). 235–251
- Patterson III, William A.** 187–200
- Pedicularis lanceolata*** 125–159
- Pennsylvania, old-growth forest** 47–63
- Peucedanum palustre*** 516–518
- Plant species associated with a regionally rare hemiparasitic plant, *Pedicularis lanceolata* (Orobanchaceae), throughout its geographic range.** 125–159; species list 141–159
- Plant-community variability in a Wisconsin calcareous fen. Historical land use and,** 160–186
- Plant-soil relations** 260–279
- Platanthera praecox*** 201–219
- Pollen** 235–251
- Pollen morphology and its relationship to taxonomy of the genus *Sarracenia* (Sarraceniaceae).** 235–251
- Pollination** 64–86; 201–219
- Pollinator effectiveness** 64–86
- Pollinator effectiveness of insect floral visitors to two sympatric species of wild aster: *Sympotrichum lanceolatum* (Willd.) Nesom and *S. lateriflorum* (L.) Löve & Löve (Asteraceae: Astereae). Relative,** 64–86
- Preliminary study of the vegetation of vernal pools of Acadia National Park, Maine, U.S.A. A,** 260–279; species list 267–273
- Primulaceae** 280–299
- Pringle, James S.** Validation of the name *Zeltnera texensis* (Gentianaceae). 514–515 (Note)
- Rajakaruna, Nishanta** 1–31; 260–279
- Rajakaruna, Nishanta, Tanner B. Harris, Stephen R. Clayden, Alison C. Dibble, and Fred C. Olday.** Lichens of the Callahan Mine, a copper- and zinc-enriched superfund site in Brooksville, Maine, U.S.A. 1–31; species list 17–31
- Rare species, *Utricularia resupinata*** 32–46; regionally rare *Pedicularis lanceolata* 125–159
- Record, Sydne.** Plant species associated with a regionally rare hemiparasitic plant, *Pedicularis lanceolata* (Orobanchaceae), throughout its geographic range. 125–159 species list 141–159
- Regionally rare plants** 125–159
- Relative pollinator effectiveness of insect floral visitors to two sympatric species of wild aster: *Sympotrichum lanceolatum* (Willd.) Nesom and *S. lateriflorum* (L.) Löve & Löve (Asteraceae: Astereae).** 64–86
- Remediation** 1–31
- Reviewers of manuscripts** 532
- Riparian forest** 47–63
- Salvia glutinosa* (Lamiaceae) naturalized in southeastern New York.** 220–224 (Note)
- Sarracenia*** 235–251
- Sarraceniaceae** 235–251

- Scribailo, Robin, W., Mitchell S. Alix, and Scott A. Namestnik. Historical notes and new records for the rare Atlantic Coastal Plain species *Utricularia resupinata* (Lentibulariaceae) in Indiana. 32-46
- Scrosati, Ricardo A. 87-105
- Seasonal variation in the nectar sugar concentration and nectar quantity in the Western Prairie Fringed Orchid, *Platanthera praecox* (Orchidaceae). 201-219
- Seaweed ecology 300-350
- Second-growth forest 47-63
- Semple, J. C. 64-86
- Serpentine 1-31
- Simon, Matt 160-186
- Sphingidae 201-219
- Spongomorpha* 87-105
- Statement of ownership 124
- Submerged fucoid algae 300-350
- Superfund site in Brooksville, Maine, U.S.A. 1-31
- Sussmann, Andrea V. and Ricardo A. Scrosati. Morphological variation in *Acrosiphonia arcta* (Codiolales, Chlorophyta) from environmentally different habitats in Nova Scotia, Canada. 87-105
- Sympyotrichum lanceolatum*, *S. lateriflorum* 64-86
- Systematics 235-251
- Tabak, Nava M. *Salvia glutinosa* (Lamiaceae) naturalized in southeastern New York. 220-224 (Note)
- Tall Grass Prairie Preserve, Manitoba 201-219
- Taxonomy 252-259
- Trichophorum planifolium* 351-352
- Tsuga canadensis* 47-63
- United States 252-259
- Utricularia resupinata* in Indiana 32-46
- Validation of the name *Zeltnera texensis* (Gentianaceae). 514-515 (Note)
- Vascular flora of Lauderdale County, Mississippi, U.S.A. The, 365-418; species list 388-418
- Vegetation of vernal pools of Acadia National Park, Maine, U.S.A. A preliminary study of the, 260-279; species list 267-273
- Vernal pool vegetation 260-279
- Weedy species 160-186
- Western Prairie Fringed Orchid, *Platanthera praecox* (Orchidaceae). 201-219
- Westwood, A. Richard, Christie L. Borkowsky, and Karen E. Budnick. Seasonal variation in the nectar sugar concentration and nectar quantity in the Western Prairie Fringed Orchid, *Platanthera praecox* (Orchidaceae). 201-219
- Williams, Charles E. 47-63
- Wisconsin, calcareous fen 160-186
- Zeltnera texensis* 514-515